



Bear Lake 2024 Aquatic Vegetation, Water Quality, and 2025 Management Recommendations Report



November, 2024

Bear Lake 2024 Aquatic Vegetation, Water Quality, and 2025 Management Recommendations Report



© Restorative Lake Sciences
Website: <http://www.restorativelakesciences.com>

Table of Contents

Section 1: Bear Lake Summary (2024).....	4
Section 2: Bear Lake Water Quality Data (Historic-2024).....	5
Section 3: Bear Lake Aquatic Vegetation Data (2024).....	15
Section 4: Bear Lake Macroinvertebrates (2024).....	20
Section 5: Management Recommendations for 2025.....	23

Bear Lake 2024 Aquatic Vegetation, Water Quality, and 2025 Management Recommendations Report

The following information is a summary of key lake findings collected in 2024.

The overall condition of Bear Lake is ranked in the top 15% of developed lakes of similar size in the state of Michigan (based on RLS evaluations). The water clarity in 2024 was good with an average of 9.3 feet. The nutrient concentrations in the lake are considered low to moderate yet are adequate to support robust submersed aquatic plant growth due to the high-water clarity. Invasive species such as Eurasian Watermilfoil (EWM) are able to grow in moderate nutrient waters and thus are a challenge to the Bear Lake ecosystem. Bear Lake had a total of 7.8 acres of EWM in 2024 (determined by whole-lake surveys conducted in May and August of 2024 and additional site surveys) which was very effectively treated with the new systemic herbicide ProcellaCOR® combined with the contact herbicide diquat just after the surveys. Rotating active ingredients is key to minimizing plant resistances. Protection of the 24 native aquatic plant species is paramount for the health of the lake fishery and these plants should not be managed unless they are a nuisance to lakefront property owners and possess navigational and recreational hazards

The lake did experience depletion of dissolved oxygen with depth during the mid-summer sampling in the East Basin, otherwise oxygen levels held consistent which is common for a large lake that minimally stratifies. Trends for most water quality parameters are shown below. Overall, Bear Lake is a very stable aquatic ecosystem with excellent aquatic vegetation biodiversity, relative abundance, and water clarity, and neutral pH and soft water alkalinity, and low to moderate nutrients.

Bear Lake Water Quality Data

Water Quality Parameters Measured

There are hundreds of water quality parameters one can measure on an inland lake, but several are the most critical indicators of lake health. These parameters include water temperature (measured in °C), dissolved oxygen (measured in mg/L), pH (measured in standard units-SU), conductivity (measured in micro-Siemens per centimeter- $\mu\text{S}/\text{cm}$), total alkalinity or hardness (measured in mg of calcium carbonate per liter-mg CaCO_3/L), total dissolved solids (mg/L), Secchi transparency (feet), total phosphorus and total nitrogen (both in mg/L), chlorophyll-*a* (in $\mu\text{g}/\text{L}$), and algal community composition. Graphs that show critical trends for some parameters in spring and late summer of each year are displayed below. Trend data were calculated using mean values for each parameter for spring and summer over each sampling location. Table 1 below demonstrates how lakes are classified based on key parameters. Bear Lake would be considered mesotrophic (relatively productive) since it does contain ample phosphorus, nitrogen, and aquatic vegetation growth but has excellent water clarity and moderate algal growth. 2024 water quality data for Bear Lake is shown below in Tables 2-4. Three deep basins were measured, with the West Basin located at the NW corner of the lake, the Central Basin located at the central portion of the lake, and the East Basin located at the northeastern portion of the lake (Figure 1). These parameters are discussed below along with water quality data specific to Bear Lake which were collected on August 21, 2024.



Table 1. Lake trophic classification (MDNR).

<i>Lake Trophic Status</i>	<i>Total Phosphorus ($\mu\text{g L}^{-1}$)</i>	<i>Chlorophyll-a ($\mu\text{g L}^{-1}$)</i>	<i>Secchi Transparency (feet)</i>
Oligotrophic	< 10.0	< 2.2	> 15.0
Mesotrophic	10.0 – 20.0	2.2 – 6.0	7.5 – 15.0
Eutrophic	> 20.0	> 6.0	< 7.5

Table 2. Bear Lake Water Quality Parameter Data Collected over the East Deep Basin on August 1, 2024.

<i>Depth ft.</i>	<i>Water Temp °C</i>	<i>DO mg L⁻¹</i>	<i>pH S.U.</i>	<i>Cond. μS cm⁻¹</i>	<i>Turb. NTU</i>	<i>TDS mg L⁻¹</i>	<i>Total Kjeldahl Nitrogen mg L⁻¹</i>	<i>Total Alk. mgL⁻¹ CaCO₃</i>	<i>Total Phos. mg L⁻¹</i>
0	27.02	8.60	8.6	230.5	0.4	147.5	<0.50	56	<0.010
12	25.07	8.72	8.6	229.1	0.7	146.6	<0.50	56	<0.010
23	23.42	3.70	7.21	262.9	1.2	168.5	0.57	55	<0.010

Table 3. Bear Lake Water Quality Parameter Data Collected over the Central Deep Basin on August 1, 2024.

<i>Depth ft.</i>	<i>Water Temp °C</i>	<i>DO mg L⁻¹</i>	<i>pH S.U.</i>	<i>Cond. μS cm⁻¹</i>	<i>Turb. NTU</i>	<i>TDS mg L⁻¹</i>	<i>Total Kjeldahl Nitrogen mg L⁻¹</i>	<i>Total Alk. mgL⁻¹ CaCO₃</i>	<i>Total Phos. mg L⁻¹</i>
0	26.67	8.32	8.58	230.4	0.9	147.4	0.5	54	<0.010
8	25.64	8.82	8.61	229.3	0.9	146.7	<0.50	54	<0.010
19	24.82	8.48	8.46	230.2	1.8	147.3	<0.50	55	<0.010

Table 4. Bear Lake Water Quality Parameter Data Collected over the West Basin on August 1, 2024.

<i>Depth ft.</i>	<i>Water Temp °C</i>	<i>DO mg L⁻¹</i>	<i>pH S.U.</i>	<i>Cond. μS cm⁻¹</i>	<i>Turb. NTU</i>	<i>TDS mg L⁻¹</i>	<i>Total Kjeldahl Nitrogen mg L⁻¹</i>	<i>Total Alk. mgL⁻¹ CaCO₃</i>	<i>Total Phos. mg L⁻¹</i>
0	26.68	8.39	8.62	230.4	0.4	147.4	<0.50	55	<0.010
10	25.57	8.74	8.58	230	0.5	147.2	0.55	54	<0.010
20	24.88	8.39	8.40	230.4	1.2	147.5	0.55	52	0.010



Figure 1: 2024 Water quality sampling locations on Bear Lake.

Dissolved Oxygen

Dissolved oxygen is a measure of the amount of oxygen that exists in the water column. In general, dissolved oxygen levels should be greater than 5 mg L^{-1} to sustain a healthy warm-water fishery. Dissolved oxygen concentrations in Bear Lake may decline if there is a high biochemical oxygen demand (BOD) where organismal consumption of oxygen is high due to respiration. Dissolved oxygen is generally higher in colder waters. Dissolved oxygen is measured in milligrams per liter (mg L^{-1}) with the use of a dissolved oxygen meter and/or through the use of Winkler titration methods. The dissolved oxygen concentrations in Bear Lake were healthy and declined minimally with depth. This may be due to an early fall turn over and reduced stratification due to constant mixing of the lake water. **The dissolved oxygen concentrations in Bear Lake support both a warm water and cool water fishery with an average concentration of around 8.1 mg L^{-1} in 2024.** Figure 2 below shows the changes in mean dissolved oxygen with time in Bear Lake.

Figure 2. Temporal Trends in Mean DO among Bear Lake Deep Basins




Figure 2: Temporal Trends in Mean Dissolved Oxygen in Bear Lake.

Water Clarity (Transparency) Data & Turbidity

Secchi transparency readings allow us to estimate where the photic to aphotic zones transition. The transparency throughout Bear Lake is adequate (mean of 9.3 feet in 2024; Figure 3) to allow abundant growth of algae and aquatic plants in the majority of the littoral zone of the lake.

Secchi transparency is variable and depends on the number of suspended particles in the water (often due to windy conditions of lake water mixing) and the amount of sunlight present at the time of measurement. Other parameters such as turbidity (measured in NTU’s) and Total Dissolved Solids (measured in mg/L) are correlated with water clarity and show an increase as clarity decreases. **The turbidity and total dissolved solids in Bear Lake during the 2024 sampling events were also quite low at ≤ 1.8 NTU’s and 149.6 mg/L, respectively.**

Total Kjeldahl Nitrogen (TKN) is the sum of organic nitrogen plus ammonia (NH₃) forms in freshwater systems. Much nitrogen (amino acids and proteins) also comprises the bulk of living organisms in an aquatic ecosystem. Nitrogen originates from atmospheric inputs (i.e., burning of fossil fuels), wastewater sources from developed areas (i.e., runoff from fertilized lawns), agricultural lands, septic systems, and from waterfowl droppings. It also enters lakes through groundwater or surface drainage, drainage from marshes and wetlands, or from precipitation (Wetzel, 2001). In lakes with an abundance of nitrogen (N: P > 15), phosphorus may be the limiting nutrient for phytoplankton and aquatic macrophyte growth. Alternatively, in lakes with low nitrogen concentrations (and relatively high phosphorus), the blue-green algae populations may increase due to the ability to fix nitrogen gas from atmospheric inputs.

Lakes with a mean TKN value of 0.66 mg L⁻¹ may be classified as oligotrophic, those with a mean TKN value of 0.75 mg L⁻¹ may be classified as mesotrophic, and those with a mean TKN value greater than 1.88 mg L⁻¹ may be classified as eutrophic. **Bear Lake possessed TKN concentrations between 0.5-0.6 mg L⁻¹ during the 2024 sampling events which is considered oligo-mesotrophic and is desirable.**

Figure 4. Temporal Trend in Mean TP among Bear Lake Deep Basins




Figure 4: Temporal trend in mean Total Phosphorous levels within Bear Lake.

Total Alkalinity

Lakes with high alkalinity ($> 150 \text{ mg L}^{-1}$ of CaCO_3) are able to tolerate larger acid inputs with less change in water column pH. Many Michigan lakes contain high concentrations of CaCO_3 and are categorized as having “hard” water.

Total alkalinity may change on a daily basis due to the re-suspension of sedimentary deposits in the water and respond to seasonal changes due to the cyclic turnover of the lake water. **The alkalinity in Bear Lake averaged 55 mg L^{-1} of CaCO_3 (Figure 5) in 2024, which is low and indicates a soft water lake.**

Figure 5. Temporal Trends in Mean Total Alkalinity among Bear Lake Deep Basins




Figure 5: Temporal trends in mean total alkalinity within Bear Lake.

pH

Most Michigan lakes have pH values that range from 6.5 to 9.5. Acidic lakes ($\text{pH} < 7$) are rare in Michigan and are most sensitive to inputs of acidic substances due to a low acid neutralizing capacity (ANC). Bear Lake is considered “slightly basic” on the pH scale. **The mean pH of Bear Lake during the 2024 sampling events averaged 8.4 S.U. which is ideal for an inland soft water lake (Figure 6).** pH values may change over time due to photosynthesis of submersed aquatic vegetation and algae.

Figure 6. Temporal Trends in Mean pH among Bear Lake Deep Basins




Figure 6: Temporal trends in mean pH within Bear Lake.

Conductivity

Conductivity is a measure of the number of mineral ions present in the water, especially those of salts and other dissolved inorganic substances. Conductivity generally increases as the amount of dissolved minerals and salts in a lake increases, and also increases as water temperature increases. **The conductivity values for Bear Lake during the 2024 sampling events (Figure 7) averaged 233.7 $\mu\text{S}/\text{cm}$ which is favorable for an inland lake.** Severe water quality impairments do not occur until values exceed 800 $\mu\text{S}/\text{cm}$ and are toxic to aquatic life around 1,000 $\mu\text{S}/\text{cm}$.

Figure 7. Temporal Trends in Mean Conductivity among Bear Lake Deep Basins




Figure 7: Temporal trends in mean Conductivity within Bear Lake.

Chlorophyll-*a* and Algal Species Composition

Chlorophyll-*a* is a measure of the amount of green plant pigment present in the water, often in the form of planktonic algae. High chlorophyll-*a* concentrations are indicative of nutrient-enriched lakes. Chlorophyll-*a* concentrations greater than 6 $\mu\text{g L}^{-1}$ are found in eutrophic or nutrient-enriched aquatic systems, whereas chlorophyll-*a* concentrations less than 2.2 $\mu\text{g/L}$ are found in nutrient-poor or oligotrophic lakes. **The mean chlorophyll-*a* concentrations during the 2024 sampling events in Bear Lake was 2.0 $\mu\text{g/L}$ which is moderately low for an inland Michigan lake and is expected given less runoff from rainfall events in 2024.**

The algal genera were determined from composite water samples collected over the deep basins of Bear Lake in 2024 were analyzed with a compound bright field microscope. The genera present included the Chlorophyta (green algae): *Scenedesmus* sp., *Chlorella* sp., *Ulothrix* sp., *Pediastrum* sp., *Cosmarium* sp., *Quadrigula* sp., *Ankistrodesmus* sp., and *Haematococcus* sp; the Bascillariophyta (diatoms): *Navicula* sp., *Fragillaria* sp., *Synedra* sp., *Stephanodiscus* sp., and *Cymbella* sp. The aforementioned species indicate a diverse algal flora and represent a good diversity of alga with an abundance of diatoms that are indicative of great water quality. Microscopic photos of some key algae in Bear Lake are shown below (Figures 8 – 11).



Figure 8: Ulothrix – A Green Algae in Bear Lake




Figure 9: *Navicula*- A diatom algae in Bear Lake




Figure 10: Pediastrum – Green Algae in Bear lake




Figure 11: *Synedra*- A diatom algae in Bear Lake

Aquatic Vegetation Data (2024)

Status of Native Aquatic Vegetation in Bear Lake

The native aquatic vegetation present in Bear Lake is essential for the overall health of the lake and the support of the lake fishery. The whole-lake species surveys in August, of 2024 determined that there were a total of 24 native aquatic plant species in Bear Lake. These included 17 submersed species, 3 floating-leaved species, and 4 emergent species. This indicates a high biodiversity of aquatic vegetation in Bear Lake. The overall % cover of the lake by native aquatic plants is moderate relative to the lake size and thus these plants should be protected unless growing near swim areas at nuisance levels. The most common native aquatic plants included: 1) Fern-leaf pondweed (Figure 12) which is a pondweed with flat leaves that tends to grow in shallow to medium depth waters. This plant resembles a small fern that is brownish-green in coloration, 2) Variable Leaf Pondweed (Figure 13) which creates vegetation beds that is a preferred habitat for fish species and macroinvertebrates. It has small, curled leaves subsurface and an oval shaped leaf on the water's surface, and 3) Illinois Pondweed (Figure 14) which has slightly broader leaves than variable leaf pondweed and also creates a canopy in the water providing fish and macro-invertebrate habitat. A list of all native aquatic plants (scientific and common names) and their relative abundance in 2024 is found in Table 5 below.




Figure 12. Fern-leaf Pondweed



Figure 13. Variable-leaf Pondweed



Figure 14. Illinois Pondweed




Figure 15: Bear Lake biovolume map from the August 1st, 2024 survey. Most of the vegetation is located in the west basin of the lake in 3-12 feet depths. Blue = no vegetation, Red = Dense vegetation.

Table 5: 2024 Bear Lake Native Aquatic Plant Species and Relative Abundance.

<i>Native Aquatic Plant Species</i>	<i>Aquatic Plant Common Name</i>	<i>% cover in/around Bear Lake (August 2024)</i>
<i>Chara vulgaris</i>	Muskgrass	17.17
<i>Potamogeton illinoensis</i>	Illinois Pondweed	31.47
<i>Potamogeton pusillus</i>	Small-leaf Pondweed	3.43
<i>Potamogeton robbinsii</i>	Fern-leaf Pondweed	45.35
<i>Stuckenia pectinata</i>	Sago Pondweed	0.14
<i>Potamogeton amplifolius</i>	Large-leaf Pondweed	4.29
<i>Potamogeton praelongus</i>	White-stem Pondweed	7.15
<i>Potamogeton gramineus</i>	Variable-leaf Pondweed	32.9
<i>Potamogeton sp.</i>	Thin Leaf Pondweed	2.72
<i>Potamogeton zosteriformis</i>	Flat-stem Pondweed	24.75
<i>Vallisneria americana</i>	Wild Celery	5.15
<i>Najas guadalupensis</i>	Southern Naiad	0.29
<i>Potamogeton perfoliatus</i>	Clasping-leaf Pondweed	0.14
<i>Megalodonta beckii</i>	Water Marigold	6.29
<i>Najas flexilis</i>	Slender Naiad	23.89
<i>Elodea canadensis</i>	Common Elodea	10.16
<i>Utricularia vulgaris</i>	Common Bladderwort	0.86
<i>Nymphaea odorata</i>	White Waterlily	0.72
<i>Nuphar variegata</i>	Yellow Waterlily	0.86
<i>Brasenia schreberi</i>	Watershield	0.29
<i>Typha latifolia</i>	Cattails	0.43
<i>Schoenoplectus acutus</i>	Bulrushes	1.29
<i>Iris versicolor</i>	Blue-flag Iris	0.14
<i>Decodon verticillatus</i>	Swamp Loosestrife	0.72

Status of Invasive (Exotic) Aquatic Plant Species in Bear Lake

The amount of Eurasian Watermilfoil (Figure 16) present in Bear Lake varies each year and is dependent upon climatic conditions, especially runoff-associated nutrients. In 2024, the EWM was low in the spring with 7.2 acres found on May 14, 2024 (Figure 17). These areas were treated by PLM with oversight by RLS the following week. Treatment utilized a combination of ProcellaCOR® and diquat. A very minor late season milfoil growth was noted during the survey on August 1, 2024 with an additional 0.7 acres of EWM (Figure 18) which was also treated shortly after with ProcellaCOR® and diquat combination.



Figure 16. Eurasian Watermilfoil



Figure 17. Distribution of EWM in Bear Lake (May 14, 2024).




Figure 18. Distribution of EWM in Bear Lake (August 1, 2024).

Bear Lake Sediment Macroinvertebrate Data (2024)

Freshwater macroinvertebrates are ubiquitous, as even the most impacted lake contains some representatives of this diverse and ecologically important group of organisms. Benthic macroinvertebrates are key components of lake food webs both in terms of total biomass and in the important ecological role that they play in processing of energy. Others are important predators, graze algae on rocks and logs, and are important food sources (biomass) for fish. The removal of macroinvertebrates has been shown to impact fish populations and total species richness of an entire lake or stream food web (Lenat and Barbour 1994). In the food webs of lakes, benthic macroinvertebrates have an intermediate position between primary producers and higher trophic levels (as fish) on the other side. Hence, they play an essential role in key ecosystem processes (food chain dynamics, productivity, nutrient cycling, and decomposition). These may also include many rare species.

Several characteristics of benthic macroinvertebrates make them useful bio-indicators of lake water quality including that many are sensitive to changes in physical, chemical, and biological conditions of a lake, many complete their life cycle in a single year, their life cycles and ecological requirements are generally well known, they are sessile organisms and cannot readily escape pollution or other negative aspect, and they are easily collected. As benthic macroinvertebrates respond sensitively not only to pollution, but also to a number of other human impacts (hydrological, climatological, morphological, navigational, recreational, and others), they could potentially be used for a holistic indication system for lake ecosystem health (Solimini et al. 2006).

Restorative Lake Sciences, LLC, collected aquatic macroinvertebrate samples from three separate locations within Bear Lake, Manistee County, Michigan on October 14, 2024 (Table 6). The sampling found five key taxa which included mayflies (*Hexagenia limbata*, Ephemeroidea), midges (Chironomidae), damselfly larvae, caddisfly larvae, and zebra mussels (Dreissenidae). Of all the species found, all were native except for the zebra mussels. While the majority of the species were native, some are located universally in low quality and high-quality water. The midge larvae family Chironomidae can be found in both high- and low-quality water (Lenat and Barbour 1994). The mayfly, *Hexagenia limbata*, found within this lake, has been shown to be linked with good water quality.

Eurasian watermilfoil (*Myriophyllum spicatum*) has been shown to negatively influence both fish and macroinvertebrate communities (Lillie and Budd 1992). In addition to exotic and invasive macroinvertebrate species, macroinvertebrate assemblages can be affected by land-use. Stewart et al. (2000) showed that macroinvertebrates were negatively affected by surrounding land-use. They also indicated that noted these land-use practices are important to restoration and management of lakes. Schreiber et al. (2003) stated that disturbance and anthropogenic land use changes are usually considered to be key factors facilitating biological invasions.

Table 6. 2024 Bear Lake Macroinvertebrates (October 14, 2024).

Sample 1 Bear Lake Marina	Sample type – Dip Net/Sediment Grab				
		Odonata	Coregonidae	2	Damselfly larvae
		Trichoptera	Limnephilidae	1	Caddis larvae
		Veneroida	Dreissenidae	5	Zebra Mussels
		Gastropoda	Physidae	11	Pond snails
			Total	19	
Sample 2 Northwest Cove	Sample type – Dip Net/Sediment Grab	Odonata	Coregonidae	2	Damselfly larvae
		Diptera	Chironomidae	2	Midge larvae
		Veneroida	Dreissenidae	9	Zebra Mussels
		Gastropoda	Physidae	8	Pond snails
			Total	21	
Sample 3 Mid-lake sample milfoil bed	Sample type – Sediment Grab				
		Odonata	Coregonidae	1	Damselfly larvae
		Gastropoda	Physidae	11	Pond snails
		Diptera	Chironomidae	5	Midge Larvae
			Total	17	

Management Recommendations for 2025

Public hearings were held on July 12, 2022 by the Bear Lake Improvement Board (BLIB) to determine if ongoing management of Bear Lake was practical. There was overwhelming support at the hearings based on feedback and public comment from Bear Lake riparians. The cost to manage Bear Lake has declined significantly over the past decade due to effective and intensive management efforts, especially regarding whole-lake species inventories. RLS cautioned the BLIB and public that at any time a seedbank re-emergence can occur in any year due to environmental conditions, and vigorous and costly treatments may be needed when this occurs. It is not usually a frequent occurrence, but all boards should have a robust reserve in case of this natural event.

In 2025, continuous whole-lake geo-referenced aquatic vegetation surveys are needed to determine the precise locations of EWM or other problematic invasives in and around Bear Lake. This is especially important since the highly invasive submersed aquatic plant Hydrilla has been found in a few Michigan ponds and numerous Midwest lakes. These surveys should occur in late May, early June, and again post-treatment in 2025 and in-between as needed if more treatments become necessary. As in previous years, scientists from RLS will be present to oversee aquatic herbicide applications.

Due to the relative scarcity of native aquatic vegetation in Bear Lake (relative to the lake area), the treatment of these species with aquatic herbicides is not recommended. The plan for 2025 includes the use of high dose systemic aquatic herbicides as used in 2024, which includes ProcettaCOR® with the contact herbicide diquat. This was used to reduce the tolerance of EWM to commonly used triclopyr and 2,4-D. The latter products may be used in 2025 for chemical product rotation to reduce herbicide tolerance by EWM if needed.

Water quality parameters in the main lake will also be monitored and graphed with historical data to observe long-term trends. This has been important since the nutrient concentrations and other parameters have demonstrated that Bear Lake is a stable and healthy aquatic ecosystem and that nutrients appear to be in balance. Nutrients are likely to increase in summer as aquatic plants grow and decay and this process results in nutrient release or during periods of heavy runoff.

Bear Lake was initially treated for over 330 acres of EWM in 2008. Since then, the lake has been treated for anywhere from 7 to 75 acres on an annual basis.

The scientists at RLS think that lake management which maintains EWM levels below 10% of the lakes surface areas are being managed successfully. Though EWM is an invasive plant that we want to minimize, it is a plant that is excellent fish cover in limited amounts. RLS surveys are intensive and site-specific, but it is impossible to find every hidden stem of EWM. RLS uses the latest technologies including GPS location of EWM and BioBase® scanning to help quantify the bio-volume of all aquatic vegetation. Below is a history of the annual acres treated to date.

<u>Year</u>	<u>Acres treated (rounded)</u>
2008	330
2009	11
2010	7
2011	11
2012	70
2013	10
2014	10
2015	29
2016	11
2017	17
2018	32
2019	75
2020	10
2021	11
2022	51
2023	24.5
2024	7.8

In conclusion, Bear Lake is a very healthy lake with excellent aquatic plant biodiversity, very good water clarity, moderate nutrients, and a healthy lake fishery. Management of the EWM and protection of the water quality are paramount for the long-term health of the lake.